



Certificate of Analysis

International Food Europe SRL
 Viale Giulio Cesare 92
 Roma 00192
 Attention: Francesco De Santis
 Phone: 800 180 766
 Email: info@naturalepiu.it

Lab Reference: 17-27787
 Submitted by:
 Date Received: 14/11/2017
 Date Completed: 16/11/2017
 Order Number:
 Reference: NaturalePiu

Report Comments

Samples were received by Analytica Laboratories in acceptable condition unless otherwise noted on this report.

Amended report. Contact details and reference changed at request of the client.

Results Summary

3in1 Honey Analysis

Laboratory ID	Sample ID	Dihydroxyacetone (DHA)	Methylglyoxal (MG)	Non-peroxide Activity* (NPA)	Hydroxymethylfurfural (HMF)
		<i>Units</i> Reporting Limit			
17-27787-3	17R9	mg/kg 10	mg/kg 4	%w/v phenol eq. 0.8	mg/kg 1
		1,570	525	15.2	9

3in1 Honey Analysis Approver:

Chris Wakefield, B.Sc.(Tech)
 Honey Team Leader

Method Summary

3in1 Determination of Dihydroxyacetone (DHA), Methylglyoxal (MG) and Hydroxymethylfurfural (HMF) by aqueous extraction, derivatisation, and UPLC analysis.

NPA Non-Peroxide Activity (NPA) values are not directly measured by the laboratory, but are calculated from the measured methylglyoxal concentration in the honey according to the requirements of the client. The calculation is based on published data^(†) comparing the NPA and methylglyoxal concentration measured in a range of honey samples. These calculated values are not accredited by IANZ and do not imply that the honey is or is not manuka honey. NPA values less than 5 are an estimate based on extrapolation of the relationship between methylglyoxal and NPA

(†) Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. And, Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey" [Carbohydr. Res. 343 (2008) 651]. Carbohydrate Research 344 (2009) 2609. C. J. Adams, et al.